

# <sup>180</sup>Ta production in the classical s-process

Markus Loewe<sup>1</sup>, Petr Alexa<sup>2</sup>, Jorrit de Boer<sup>1</sup>, Michael Würkner<sup>1</sup>

<sup>1</sup>*Sektion Physik, Munich University, Am Coulombwall 1, Garching, D-85768, Germany*

<sup>2</sup>*Department of Physics and Measurements, Institute of Chemical Technology,  
Technická 5, CZ-166 28 Prague 6, Czech Republic*

(February 1, 2008)

The production and survival of the quasistable isomer <sup>180</sup>Ta during the stellar nucleosynthesis has remained a matter of discussion for years. A careful analysis of the available experimental data and theoretical calculations enabled us to reproduce the observed solar abundance of <sup>180</sup>Ta even in the classical s-process ( $kT = 28 \text{ keV} - 33 \text{ keV}$ ).

26.20.+f, 26.30.+k, 27.70.+q, 97.10.Cv

<sup>180</sup>Ta is the only nucleus present in nature in an  $9^-$  isomeric state (<sup>180</sup>Ta<sup>m</sup>) at an energy of 75.3 keV. For its half-life an experimental lower limit of  $1.2 \times 10^{15} \text{ y}$  was obtained [1]. The  $1^+$  ground state of <sup>180</sup>Ta decays to <sup>180</sup>Hf and <sup>180</sup>W (half-life 8.15 h). <sup>180</sup>Ta<sup>m</sup> has a very small abundance: Only 0.012% of natural Ta consists of <sup>180</sup>Ta<sup>m</sup>, the rest is <sup>181</sup>Ta. The solar abundance of <sup>180</sup>Ta<sup>m</sup> (normalized to Si with the solar abundance of  $10^{12}$ ) is 2.48 (cf. solar abundances of neighbouring isotopes:  $4.20 \cdot 10^4$  for <sup>178</sup>Hf,  $2.10 \cdot 10^4$  for <sup>179</sup>Hf,  $5.4 \cdot 10^4$  for <sup>180</sup>Hf, and 173 for <sup>180</sup>W) [2].

To explain the production of <sup>180</sup>Ta<sup>m</sup> during the nucleosynthesis, two types of production processes were proposed.

## I. Explosive processes occur in supernovae:

1. The r-process leading to an isomer in <sup>180</sup>Lu decaying to an  $8^-$  isomer in <sup>180</sup>Hf at 1.14 MeV, further decaying partially to <sup>180</sup>Ta<sup>m</sup> [3,4]: The amount of <sup>180</sup>Ta<sup>m</sup> produced in the r-process relative to the amount of <sup>180</sup>Hf can be obtained from [5]:

$$\frac{N_r(^{180}\text{Ta}^m)}{N_r(^{180}\text{Hf})} = f_m^{180} \cdot f_{\beta^-}^m, \quad (1)$$

where  $f_m^{180}$  and  $f_{\beta^-}^m$  are the branching factors for the  $\beta^-$  or  $\beta^- + \gamma$  transitions  $^{180}\text{Lu} \rightarrow ^{180}\text{Hf}^m$  and  $^{180}\text{Hf}^m \rightarrow ^{180}\text{Ta}^m$ , respectively. For  $f_{\beta^-}^m$  a value of  $(0.29 \pm 0.08)\%$  was found [4]. The laboratory value can also be used for the r-process. Corrections due to the high degree of ionization of <sup>180</sup>Hf<sup>m</sup> can be neglected since the r-process duration ( $\leq 100 \text{ s}$ ) is short compared to the isomer half-life (5.5 hr) and the temperature and density are greatly diminished when the decay of <sup>180</sup>Hf<sup>m</sup> occurs. Values of  $(0.46 \pm 0.15)\%$  [3] or  $(0.005 \pm 0.018)\%$  [6] were found for  $f_m^{180}$ . These two values of  $f_m^{180}$  lead to  $(9 \pm 4)\%$  of the observed <sup>180</sup>Ta<sup>m</sup> solar abundance or to a negligible <sup>180</sup>Ta<sup>m</sup> production.

2. In the rapid p-process, <sup>180</sup>Ta<sup>m</sup> can be produced in the <sup>181</sup>Ta( $\gamma, n$ ) reaction (a negligible production [7] or an overproduction comparable to that of neighboring nuclei [8]).
3. In the  $\nu$ -process during the supernova-core collapse into a neutron star, <sup>180</sup>Ta<sup>m</sup> can be produced in the <sup>181</sup>Ta( $\nu, \nu' n$ ) reaction [9,10].

## II. Non-explosive processes can be summarized as follows:

1. The s-process branching in <sup>180</sup>Hf: In the s-process via <sup>179</sup>Hf( $n, \gamma$ ), the  $8^-$  isomer in <sup>180</sup>Hf partially decaying to <sup>180</sup>Ta<sup>m</sup> (with the branching factor  $f_{\beta^-}^m$ ) is populated with the branching factor  $\text{Hf}f_n^m$  or  $B$  [11]. We assume that the classical s-process lasts more than 1 year, the temperature lies between  $kT = 28 \text{ keV}$  and  $33 \text{ keV}$  [12], the neutron density  $n_n = (4.1 \pm 0.6) \cdot 10^8 \text{ cm}^{-3}$  [13], the electron density  $n_e = 5.4 \cdot 10^{26} \text{ cm}^{-3}$  [14], and  $kT$ ,  $n_n$ , and  $n_e$  remain constant during the s-process. For the relative abundance of <sup>180</sup>Ta<sup>m</sup> one then obtains:

$$\left[ \frac{N_s(^{180}\text{Ta}^m)}{N_s(^{179}\text{Hf})} \right]_1 = \frac{\langle \sigma \rangle_{^{179}\text{Hf}+n}}{\langle \sigma \rangle_{^{180}\text{Ta}^m+n}} \cdot \text{Hf}f_n^m \cdot f_{\beta^-}^m, \quad (2)$$

where  $\langle \sigma \rangle_{^{179}\text{Hf}+n}$  and  $\langle \sigma \rangle_{^{180}\text{Ta}^m+n}$  are Maxwellian averaged neutron-capture cross sections taken from [5] (Table VIII) and [15,16] (Table 9), respectively. For  $\langle \sigma \rangle_{^{179}\text{Hf}+n} = (991 \pm 30) \text{ mb}$ ,  $\langle \sigma \rangle_{^{180}\text{Ta}^m+n} = (1465 \pm 100) \text{ mb}$  ( $kT = 30 \text{ keV}$ ),  $\text{Hf}f_n^m = \sigma^m(^{179}\text{Hf})/\sigma(^{179}\text{Hf}) = (1.24 \pm 0.06)\%$  [5], and  $f_{\beta^-}^m \approx 0.7\%$  [5], this s-process branching can account for only  $(16 \pm 3)\%$  of the observed <sup>180</sup>Ta<sup>m</sup> solar abundance.

2. The s-process branching in <sup>179</sup>Hf: Excited states in <sup>179</sup>Hf decaying to <sup>179</sup>Ta can be thermally populated. <sup>180</sup>Ta<sup>m</sup> is then produced in the <sup>179</sup>Ta( $n, \gamma$ ) reaction [17] with the branching factor  $\text{Ta}f_n^m$ . For the relative abundance of <sup>180</sup>Ta<sup>m</sup> one then obtains:

$$\left[ \frac{N_s(^{180}\text{Ta}^m)}{N_s(^{178}\text{Hf})} \right]_2 = \frac{\langle \sigma \rangle_{^{178}\text{Hf}+n}}{\langle \sigma \rangle_{^{180}\text{Ta}^m+n}} \cdot \text{Ta}f_n^m \cdot f_{180}, \quad (3)$$

where  $f_{180}$  is the branching factor for neutron captures at <sup>178</sup>Hf leading to <sup>180</sup>Ta,

$$f_{180} = \left\{ \frac{\lambda(^{179}\text{Hf}+n)}{\lambda(^{179}\text{Hf})_{\beta^-}} \cdot \left[ \frac{\lambda(^{179}\text{Ta})_{EC}}{\lambda(^{179}\text{Ta}+n)} + 1 \right] + 1 \right\}^{-1} \quad (4)$$

and  $\lambda$  are transition rates. For  $kT = 30$  keV,  $\langle\sigma\rangle_{178\text{Hf}+n} = 310$  mb [5],  $\text{Ta}_n^m \approx (4.3 \pm 0.8)\%$  [18]. The other parameters are taken from [19] (an error of  $\pm 30\%$  assumed) and interpolated for the  $n_e$  and  $T$ . For  $kT = 30$  keV this s-process branching yields  $(190 \pm 40)\%$  of the  $^{180}\text{Ta}^m$  solar abundance.

It should be noted that all  $^{180}\text{W}$  can be produced in the s-process via the decay of the  $^{180}\text{Ta}$  ground-state. For the relative abundance of  $^{180}\text{W}$  one can write

$$\frac{N_s(^{180}\text{W})}{N_s(^{178}\text{Hf})} = \frac{\langle\sigma\rangle_{178\text{Hf}+n}}{\langle\sigma\rangle_{180\text{W}+n}} \cdot (1 - \text{Ta}_n^m) \cdot f_{180} \cdot f_{\beta^-}, \quad (5)$$

where  $f_{\beta^-}$  is the branching factor for the  $^{180}\text{Ta}$  ground-state decay to  $^{180}\text{W}$  (numerical values taken from [18,19]). For  $kT = 30$  keV this process can account for  $(95 \pm 36)\%$  of the  $^{180}\text{W}$  solar abundance.

3. In the s-process during the He shell burning in the AGB phase of  $1.5$  to  $3M_\odot$  mass stars about  $85\%$  of  $^{180}\text{Ta}^m$  can be produced [16].
4. The p-process in highly evolved massive stars: During the presupernova phase under temperatures  $T > 10^9$  K, thermal photons can induce the reaction  $^{181}\text{Ta}(\gamma, n)$  populating  $^{180}\text{Ta}^m$  [20].
5.  $^{180}\text{Ta}^m$  production in the cosmic radiation: Protons from the low-energy component of the galactic cosmic radiation can produce  $^{180}\text{Ta}^m$  via the (p,yp xn) reaction on s-process or r-process nuclei in the interstellar medium [21].




FIG. 1. The total s-process abundance of  $^{180}\text{Ta}^m$  (solid line) and  $^{180}\text{W}$  (long-dashed line) calculated relative to the  $^{178}\text{Hf}$  abundance as a function of the thermal energy  $kT$  or temperature  $T$ . Error bands are depicted by dotted lines. The s-process temperature window [12] is marked by dot-dashed lines.

The total relative s-process abundance of  $^{180}\text{Ta}^m$  can be calculated from:

$$\frac{N_s(^{180}\text{Ta}^m)}{N_s(^{178}\text{Hf})} = \frac{N_s(^{179}\text{Hf})}{N_s(^{178}\text{Hf})} \cdot \left[ \frac{N_s(^{180}\text{Ta}^m)}{N_s(^{179}\text{Hf})} \right]_1 + \left[ \frac{N_s(^{180}\text{Ta}^m)}{N_s(^{178}\text{Hf})} \right]_2 \quad (6)$$

From Fig. 1 one can see that for  $kT = 30$  keV an amount of  $^{180}\text{Ta}^m$  ( $2.0 \pm 0.5$ ) times larger than the observed one can be produced in the classical s-process.




FIG. 2. Internal and external population and depopulation possibilities in the three-level system described by means of transition rates  $\lambda$  [22].

This overproduction is reduced by intermediate states (IS) in  $^{180}\text{Ta}$ : In the s-process site thermal photons may excite higher-lying levels which then decay back either to the  $1^+$  ground state or to the  $9^-$  isomer. To find these levels Belic *et al.* [10,23] used the Stuttgart Dynamitron facility with both enriched (5.6%) and natural Ta targets. Irradiations were performed for bremsstrahlung endpoint energies  $E_0 = 0.8$ – $3.1$  MeV. Depopulation of the isomer was observed down to  $E_0 \approx 1.01$  MeV. This means that the lowest IS may have an excitation energy  $E_{IS} = 1.085$  MeV (above the ground state). Assuming that  $E_{IS}$  is the excitation energy of the lowest IS, the experimental total integrated depopulation cross section  $I_D$  turns out to be  $(5.7 \pm 1.2)$  eV fm<sup>2</sup>. Then the effective lifetime of the IS for the depopulation into the ground state  $\tau_{\text{eff}}$  is roughly equal to  $6 \cdot 10^{-11}$  s.

The IS  $|IS\rangle$  decays via a  $\gamma$ -cascade, and lifetimes and energies of the decay states  $|k\rangle$  are not known. Klay [24] showed that multi-step transitions from  $|IS\rangle$  to the ground-state  $|g\rangle$  can be substituted by a direct transition from  $|IS\rangle$  to  $|g\rangle$  if the states  $|k\rangle$  possess short enough lifetimes  $\tau_k$ , i.e.

$$\tau_k \ll \tau_{\text{eff}} \cdot \frac{2I_k + 1}{2I_{IS} + 1} \cdot \exp(|E_{IS} - E_k|/kT) \quad (7)$$

where  $E_{IS}$  and  $I_{IS}$ ,  $E_k$  and  $I_k$  are the energies and spins of the IS and the states  $|k\rangle$ , respectively. It can be shown that the condition (7) is fulfilled in  $^{180}\text{Ta}$  and transitions from the isomer  $|m\rangle$  to the ground-state  $|g\rangle$  can be studied in the three-level system (see Fig. 2).

The population of the three-level system can be described by the following coupled inhomogeneous linear differential equations:

$$\begin{aligned}\frac{dN_m}{dt} &= \lambda_{pm} N_s - (\lambda_{mIS} + \lambda_{md}) N_m + \lambda_{ISm} N_{IS} \\ \frac{dN_{IS}}{dt} &= \lambda_{pIS} N_s + \lambda_{gIS} N_g + \lambda_{mIS} N_m \\ &\quad - (\lambda_{ISg} + \lambda_{ISm} + \lambda_{ISd}) N_{IS} \\ \frac{dN_g}{dt} &= \lambda_{pg} N_s - (\lambda_{gIS} + \lambda_{gd}) N_g + \lambda_{ISg} N_{IS},\end{aligned}\quad (8)$$

where  $\lambda_{px}$  ( $\lambda_{xd}$ ) are the population (depopulation) transition rates of the state  $|x\rangle$ ,  $\lambda_{xy}$  are the transition rates between the states  $|x\rangle$  and  $|y\rangle$ ,  $N_x$  is the number of nuclei in the state  $|x\rangle$ ,  $N_x(t=0) = 0$ , and  $N_s$  is a constant number of seed nuclei ( $^{178}\text{Hf}$  in our case).

In previous analyses, e.g. [23,25], the effect of an IS on the survival of  $^{180}\text{Ta}^m$  in the presence of a stellar photon bath was calculated by solving coupled differential equations for the three-level system isomer  $\leftrightarrow$  IS  $\leftrightarrow$  ground-state without taking into account the population of the three levels due to the s-process simultaneously, i.e.  $\lambda_{pm} = \lambda_{pIS} = \lambda_{pg} = 0$  was assumed.




FIG. 3. Survival ratio  $R$  of  $^{180}\text{Ta}^m$  as a function of  $kT$  or  $T$  for the IS at  $E_{IS} = 1.085$  MeV. Error bands are marked by dotted lines, the s-process temperature window by dot-dashed lines.

For  $t \rightarrow \infty$  the solution of (8) approaches the equilibrium solution. Since the maximum equilibrium relaxation time  $\tau_{max}$  ( $\tau_{max} < 100$  days for  $kT$  between 15 keV and 45 keV) is smaller than the s-process duration ( $> 1$  year) the exact solution of (8) can be approximated by the equilibrium solution. In the case of existing IS one

can define the survival ratio  $R$  of  $^{180}\text{Ta}^m$  (depicted in Fig. 3):

$$R \equiv \frac{N_m(t \rightarrow \infty)}{N_s(^{178}\text{Hf})} / \frac{N_s(^{180}\text{Ta}^m)}{N_s(^{178}\text{Hf})} \quad (9)$$

where  $N_s(^{178}\text{Hf}) = N_s$  in (8),  $N_s(^{180}\text{Ta}^m)/N_s(^{178}\text{Hf})$  represents the relative abundance for no IS and is obtained from (6). The survival ratio  $R$  is larger than 0.2 in the whole temperature interval. For  $kT \leq 21$  keV the coupling between the isomer and the ground state via the IS is negligible. Note that  $R > 1$  for  $kT \gtrsim 40$  keV.

It can be shown that for  $kT = 30$  keV only dominant transition rates  $\lambda$  must be taken into account and that the equilibrium solution can be approximated by

$$\frac{N_m(t \rightarrow \infty)}{N_s(^{178}\text{Hf})} = \left[ \frac{N_s(^{180}\text{Ta}^m)}{N_s(^{178}\text{Hf})} \right]_{IS} \approx \frac{\lambda_{pm} + \lambda_{pg} + \lambda_{pIS}}{\lambda_{gd}/P + \lambda_{md}} \quad (10)$$

where

$$P = \frac{2I_m + 1}{2I_g + 1} \cdot \exp(-75.3 \text{ keV}/kT) \approx 0.5, \quad (11)$$

$$\lambda_{md} = \lambda(^{180}\text{Ta}^m + n) = n_n \cdot \left( \frac{2kT}{m_n} \right)^{1/2} \langle \sigma \rangle_{^{180}\text{Ta}^m + n}, \quad (12)$$

$\lambda_{md} \approx 10^{-7} \text{ s}^{-1}$ ,  $m_n$  is the neutron mass,  $\lambda_{gd} = \lambda(^{180}\text{Ta})_{\beta^-} + \lambda(^{180}\text{Ta})_{EC} = 4.2 \cdot 10^{-6} \text{ s}^{-1}$  [19],

$$\lambda_{pm} + \lambda_{pg} + \lambda_{pIS} \approx \lambda(^{178}\text{Hf} + n) \cdot \left( f_{180} + {}^{\text{Hf}}f_n^m \cdot f_{\beta^-}^m \right), \quad (13)$$

where  $\lambda(^{178}\text{Hf} + n)$  can be calculated from  $\langle \sigma \rangle_{^{178}\text{Hf} + n}$ . The two terms in (13) originate in the two s-process branchings via  $^{179}\text{Hf}$  and  $^{180}\text{Hf}^m$  that can produce  $^{180}\text{Ta}$ . As a consequence of (10) the approximate solution does not depend on properties of the IS like its spin, energy, lifetime and transition rates to isomer and ground state.

In Fig. 4 the relative abundance of  $^{180}\text{Ta}^m$  as a function of  $kT$  or  $T$  is depicted. The equilibrium solution of (8) with the IS at 1.085 MeV is compared to the case of no IS (cf. Fig. 1). Note that for  $kT \geq 29$  keV the approximate (10) and the equilibrium solution are indistinguishable. As can be seen in Fig. 4 exactly 100% of the solar  $^{180}\text{Ta}^m$  abundance can be reproduced in the middle of the s-process temperature window. Other IS that may be found in the future will not change this result (cf. a possible IS below 737 keV [26]).

If the IS at 1.085 MeV in  $^{180}\text{Ta}$  is taken into account the relative abundance of  $^{180}\text{W}$  (5) must be corrected:

$$\left[ \frac{N_s(^{180}\text{W})}{N_s(^{178}\text{Hf})} \right]_{IS} \approx \frac{\langle \sigma \rangle_{^{178}\text{Hf} + n}}{\langle \sigma \rangle_{^{180}\text{W} + n}} \cdot \left( 1 - \text{Ta}f_n^m R \right) \cdot f_{180} \cdot f_{\beta^-}. \quad (14)$$

Since the survival ratio  $R \leq 1$  for the s-process temperature window (see Fig. 3) and  $T_{af_n}^m = 4.3\%$  the maximum change of the relative abundance of  $^{180}\text{W}$  due to the IS represents about 4% and can be neglected.




FIG. 4. Comparison of the total s-process abundances of  $^{180}\text{Ta}^m$  calculated relative to the  $^{178}\text{Hf}$  abundance as a function of the thermal energy  $kT$  or temperature  $T$ : for the IS at 1.085 MeV the equilibrium solution of (8) is denoted by a bold solid line, and the approximate solution (10) by a bold long-dashed line, and the total abundance for no IS from (6) by a bold dashed line. Dotted lines correspond to error bands, dot-dashed lines to the s-process temperature window.

We have shown that  $^{180}\text{Ta}^m$  could be produced in the classical s-process dominantly via the branching in  $^{179}\text{Hf}$ . The production of  $^{180}\text{Ta}^m$  is then connected to the production of  $^{180}\text{W}$ . For the temperature window  $kT = (28 - 33)$  keV we obtain  $(50 \pm 20)\% - (220 \pm 80)\%$  of  $^{180}\text{W}$  produced in the s-process via  $^{180}\text{Ta}$ . Assuming sufficiently low temperatures in the star evolution phases following the s-process and the IS at  $E_{IS} \leq 1.085$  MeV as observed in the Stuttgart photoactivation experiment [23,10], the s-process production of  $^{180}\text{Ta}^m$  ranges from  $(30 \pm 10)\%$  to  $(230 \pm 80)\%$  of its observed solar abundance for the same temperature window. For the middle of the temperature window ( $kT = 30.5$  keV) we obtain  $(90 \pm 30)\%$  of  $^{180}\text{Ta}^m$  (see Fig. 4) and  $(110 \pm 40)\%$  of  $^{180}\text{W}$ . It should be noted that  $^{180}\text{W}$  is often listed among p-process nuclei [8]. An upper limit for its production in the s-process provided by an improved model of the p-process would have direct consequences for the  $^{180}\text{Ta}^m$  production in the classical s-process via the branchings in  $^{179}\text{Hf}$  and  $^{180}\text{Hf}$ .

#### ACKNOWLEDGMENTS

The authors are grateful to Dr. F. Käppeler and D. Brandmaier for valuable discussions.

[1] J.B. Cumming, D.E. Alburger, Phys. Rev. C **31**, 1494 (1985).

- [2] E. Anders and N. Grevesse, Geochim. Cosmochim. Acta **53**, 197 (1989).
- [3] W. Eschner, W.D. Schmitt-Ott, E. Runte, L. Gippert, H. Beer, G. Walter, R. Kirchner, O. Klepper, E. Roeckl, and D. Schardt, Z. Phys. **A317**, 281 (1984).
- [4] S.E. Kellogg and E.B. Norman, Phys. Rev. C **46**, 1115 (1992).
- [5] H. Beer and R.L. Macklin, Phys. Rev. C **26**, 1404 (1982).
- [6] S.E. Kellogg and E.B. Norman, Phys. Rev. C **34**, 2248 (1986).
- [7] S.E. Woosley and W.M. Howard, Astrophys. J. Suppl. Series **36**, 285 (1978).
- [8] M. Rayet, M. Arnould, M. Hashimoto, N. Prantzos, and K. Nomoto, Astron. Astrophys. **298**, 517 (1995).
- [9] S.E. Woosley, D.H. Hartmann, R.D. Hoffman, and W.C. Haxton, Astrophys. J. **356**, 272 (1990).
- [10] D. Belic, C. Arlandini, J. Besserer, J. de Boer, J.J. Carroll, J. Enders, T. Hartmann, F. Käppeler, H. Kaiser, U. Kneissl, E. Kolbe, K. Langanke, M. Loewe, H.J. Maier, H. Maser, P. Mohr, P. von Neumann-Cosel, A. Nord, H.H. Pitz, A. Richter, M. Schumann, F.-K. Thielemann, S. Volz, A. Zilges, Phys. Rev. C **65**, 035801 (2002).
- [11] H. Beer and E. Warde, Nature (London) **291**, 308 (1981).
- [12] K. Wisshak, F. Voss, F. Käppeler, K. Guber, L. Kazakov, N. Kornilov, M. Uhl, and G. Reffo, Phys. Rev. C **52**, 2762 (1995).
- [13] K.A. Toukan, K. Debus, F. Käppeler, and G. Reffo, Phys. Rev. C **51**, 1540 (1995).
- [14] J. Best, Forschungszentrum Karlsruhe Report No. 5824 (1996).
- [15] K. Wisshak, F. Voss, C. Arlandini, F. Käppeler, M. Heil, R. Reifarh, M. Kr̃ĩčka, and F. Bẽṽar̃, Forschungszentrum Karlsruhe Report No. 6362 (2000).
- [16] K. Wisshak, F. Voss, C. Arlandini, F. Bẽṽar̃, O. Straniero, R. Gallino, M. Heil, F. Käppeler, M. Kr̃ĩčka, S. Maser, R. Reifarh, and C. Travaglio, Phys. Rev. Lett. **87**, 251102 (2001).
- [17] K. Yokoi and K. Takahashi, Nature (London) **305**, 198 (1983).
- [18] Zs. Németh, F. Käppeler, and G. Reffo, Astrophys. J. **392**, 277 (1992).
- [19] K. Takahashi and K. Yokoi, Atomic Data Nucl. Data Tables **36**, 375 (1987).
- [20] M. Arnould, Astron. Astrophys. **46**, 117 (1976).
- [21] K.L. Hainebach, D.N. Schramm, and J.B. Blake, Astrophys. J. **205**, 920 (1976).
- [22] R.A. Ward and W.A. Fowler, Astrophys. J. **238**, 266 (1980).
- [23] D. Belic, C. Arlandini, J. Besserer, J. de Boer, J.J. Carroll, J. Enders, T. Hartmann, F. Käppeler, H. Kaiser, U. Kneissl, M. Loewe, H.J. Maier, H. Maser, P. Mohr, P. von Neumann-Cosel, A. Nord, H.H. Pitz, A. Richter, M. Schumann, S. Volz, and A. Zilges, Phys. Rev. Lett. **83**, 5242 (1999).
- [24] N. Klay, PhD. Thesis, Forschungszentrum Karlsruhe Report No. 4675 (1990).
- [25] M. Schumann, PhD. Thesis, Forschungszentrum Karlsruhe Report No. 5985 (1997).
- [26] L. Lakosi, T.C. Nguyen, Nucl. Phys. A **697**, 44 (2002).